

QUANTUM COHOMOLOGY RINGS OF GRASSMANNIANS AND TOTAL POSITIVITY

KONSTANZE RIETSCH

ABSTRACT. We give a proof of a result of D. Peterson's identifying the quantum cohomology ring of a Grassmannian with the reduced coordinate ring of a certain subvariety of GL_n . The totally positive part of this subvariety is then constructed and we give closed formulas for the values of the Schubert basis elements on the totally positive points. We then use the developed methods to give a new proof of a formula of Vafa and Intriligator and Bertram for the structure constants (Gromov–Witten invariants). Finally, we use the positivity of these Gromov–Witten invariants to prove certain inequalities for Schur polynomials at roots of unity.

1. INTRODUCTION

The group U^+ of unipotent upper-triangular matrices in GL_n have on their coordinate ring a nice basis with positive structure constants. Namely one has the dual of the classical limit of Lusztig's geometrically constructed canonical basis of the quantized enveloping algebra. The existence of this basis has been closely tied [10] to the 'totally positive part' of U^+ (the matrices with only nonnegative minors).

In this paper we study certain remarkable subvarieties $\mathcal{V}_{d,n}$ of U^+ which come up in the stabilizer of a particular standard principal nilpotent element e as closures of the 1-dimensional components under Bruhat decomposition. By a theorem of Dale Peterson's [16] the quantum cohomology rings of Grassmannians may be identified with the coordinate rings of these varieties. Therefore like U^+ itself these varieties have coordinate rings with 'canonical' bases on them (this time coming from Schubert bases), and with positive structure constants.

Most of Peterson's results, in particular this one, are unpublished. But [8] is at least a reference for the "Peterson variety" which is Peterson's approach to encoding all the quantum cohomology rings of partial flag varieties in one go, and into which the varieties $\mathcal{V}_{d,n}$ may be embedded naturally.

We study the varieties $\mathcal{V}_{d,n}$ here from the point of view of explicitly describing their totally positive parts. We show that the totally positive elements come in a single one-parameter family, and give closed formulas for the values of the Schubert basis on it. Furthermore we obtain an upper bound (coming from this totally positive part) for certain Schur polynomials evaluated at roots of unity.

Date: July 25, 2000, revised September 10, 2001.

1991 Mathematics Subject Classification. 20G20, 15A48, 14N35, 14N15, 05E05.

Key words and phrases. Grassmannians, quantum cohomology, total positivity.

This work was begun at the Institute for Advanced Study with the support of NSF grant DMS 97-29992. The author is currently supported by EPSRC grant GR/M09506/01.

Since our main approach is to study the coordinate rings directly and by elementary means, we also find some new very simple proofs for results about them implied by the isomorphism with quantum cohomology. This includes properties of their Schubert bases, their explicit presentation (and thus Peterson's theorem), and finally a formula of Vafa and Intriligator and Bertram for the structure constants.

The paper is organized as follows. After an initial section on the centralizer of the principal nilpotent element e and its Bruhat decomposition we focus in on the special subvarieties $\mathcal{V}_{d,n}$ mentioned above. In Section 3 we state Theorem 3.4 on the coordinate ring of $\mathcal{V}_{d,n}$ and prove it partially. What makes the variety $\mathcal{V}_{d,n}$ so accessible to elementary methods is that its points are very easy to construct explicitly. This is explained next. In Section 4 we prove some orthogonality formulas for values of Schur polynomials at roots of unity. These facilitate the final step in the proof of Theorem 3.4 and are also the essential ingredient for the proof of the Bertram–Vafa–Intriligator formula in Section 6. Finally, after a little review of total positivity in Section 7, we come to determining the totally positive part of $\mathcal{V}_{d,n}$. The totally nonnegative matrices in $\mathcal{V}_{d,n}$ have a very beautiful explicit description, as do the values the Schubert basis elements take on them. In Section 9 we write down how these totally positive matrices factor into products of elements of simple root subgroups.

In Section 10 we explain the interpretation of the coordinate ring of $\mathcal{V}_{d,n}$ in terms of quantum cohomology and put into context some of the results encountered in the earlier sections. We also note that one property of this ring that we have not been able to derive in an elementary way and which is not at present known other than following from the geometric definitions, is the positivity of the structure constants. (The Vafa–Intriligator formula which computes these is worse than alternating). We use this positivity property in the final Section 11 to prove an inequality for values of Schur polynomials at roots of unity.

Acknowledgements. I was very fortunate to hear many inspiring lectures by Dale Peterson and would like to thank him here. I am also grateful to Shahn Majid for some useful discussions during the writing of this paper.

2. THE STABILIZER OF A PRINCIPAL NILPOTENT

2.1. Preliminaries. We recall some standard facts and notations for $GL_n(\mathbb{C})$. Let B^+, B^- be the subgroups of upper-triangular respectively lower-triangular matrices in $GL_n(\mathbb{C})$, and U^+ and U^- their unipotent radicals. Their intersection is the maximal torus T of all diagonal matrices in $GL_n(\mathbb{C})$. Let $e_1, \dots, e_{n-1}$ and $f_1, \dots, f_{n-1}$ be the standard Chevalley generators in the Lie algebras of U^+ and U^- , respectively. So e_i is the $n \times n$ -matrix with unique non-zero entry 1 found in position $(i, i+1)$ and f_i its transpose. Let $X^*(T)$ denote the character group of T , written additively, and $\Delta \subset X^*(T)$ the set of roots. Denote by $\Pi = \{\alpha_i \in \Delta \mid i = 1, \dots, n-1\}$ the usual system of simple roots defined by $te_it^{-1} = \alpha_i(t)e_i$ for all $t \in T$. The resulting sets of positive and negative roots are denoted by Δ^+ respectively Δ^- .

We identify the Weyl group of GL_n , the symmetric group, with the group $W \subset GL_n(\mathbb{C})$ of permutation matrices. And we let $s_1, \dots, s_{n-1}$ denote the usual Coxeter generators of W . So s_j corresponds to the adjacent transposition $(j, j+1)$. W acts in the usual way on $X^*(T)$ preserving Δ , and for any $w \in W$, the length $\ell(w)$ is

the number of positive roots sent to negative by w . Let w_0 be the order-reversing permutation, so the longest element in W .

2.2. $(U^+)^e$ and its Bruhat decomposition.

Definition 2.1. Let us fix the principal nilpotent element

$$e = \sum_{i=1}^{n-1} e_i \in \mathfrak{gl}_n(\mathbb{C}).$$

Let $(U^+)^e := \{u \in U^+ \mid ueu^{-1} = e\}$, the stabilizer of e in U^+ . This is an abelian subgroup of U^+ of dimension $n - 1$. The elements of $(U^+)^e$ are precisely those elements of U^+ of the form

$$(2.1) \quad u = \begin{pmatrix} 1 & x_1 & x_2 & \cdots & x_{n-2} & x_{n-1} \\ & 1 & x_1 & & & x_{n-2} \\ & & \ddots & \ddots & & \vdots \\ & & & \ddots & x_1 & x_2 \\ & & & & 1 & x_1 \\ & & & & & 1 \end{pmatrix}.$$

We can thereby explicitly identify the coordinate ring $\mathbb{C}[(U^+)^e]$ with the polynomial ring $\mathbb{C}[x_1, \dots, x_{n-1}]$.

2.3. To decompose $(U^+)^e$ by the Bruhat decomposition we need to look a bit more closely at the Weyl group. Let K be a subset of $\{1, \dots, n-1\}$. Then to K we associate the parabolic subgroup $W_K = \langle s_k \rangle_{k \in K}$ of W and its unique longest element w_K . We also consider the element $w^K := w_0 w_K$, which is the minimal length coset representative in $w_0 W_K$. Let $\{\hat{k}\}$ be the set $\{1, \dots, n-1\} \setminus \{k\}$.

We begin with two lemmas which we learned from Dale Peterson.

Lemma 2.2 ([15]). *The elements $w_K \in W$ defined above are characterized by*

$$\{w_K \in W \mid K \subset \{1, \dots, n-1\}\} = \{w \in W \mid w \cdot \Pi \subset (-\Pi) \cup \Delta^+\}.$$

Proof. Suppose that $w = w_K$ is the longest element in $W_K = \langle s_k \mid k \in K \rangle$. Then w_K has the property that it sends $\Pi_K = \{\alpha_k \mid k \in K\}$ to $-\Pi_K$, while all other simple roots remain positive. This implies the inclusion $\subseteq$.

Now let $w \in W$ lie in the right-hand side. So $w \cdot \Pi \subset (-\Pi) \cup \Delta^+$. Then let K be defined by $K = \{j \mid -\alpha_j \in w \cdot \Pi\}$. By this definition $w_K^{-1} w \cdot \Pi \subset \Delta^+$. Therefore $w = w_K$. Note that $w_K^{-1} w \cdot \alpha_i$ is positive also when $w \cdot \alpha_i \in \Delta^+$ since any positive root sent to $-\Delta^+$ by w_K must lie in $\sum_{k \in K} \mathbb{N} \alpha_k$. This cannot be true of $w \cdot \alpha_i$ since $w^{-1} \cdot \alpha_k \in -\Pi$ for any $k \in K$. $\square$

Lemma 2.3 ([15]). *Bruhat decomposition induces*

$$(2.2) \quad (U^+)^e = \bigsqcup_{K \subset \{1, \dots, n-1\}} (U^+)^e \cap B^- w^K B^-.$$

Proof. By Bruhat decomposition we can write $u = b_1 w_0 w b_2$ for some $b_1 \in B^-$, $b_2 \in U^-$ and $w \in W$. Since $u \cdot e := ueu^{-1} = e$, we have

$$w b_2 \cdot e = w_0 b_1^{-1} \cdot e.$$

The right-hand side of this equation is of the form $\sum_{j=1}^{n-1} m_j f_j + x$ for an upper-triangular matrix x and some $m_j \in \mathbb{C}$. The left-hand side is $w \cdot (e + y)$ for some lower-triangular matrix y . It follows from their equality that $w \cdot \Pi \subset (-\Pi) \cup \Delta^+$. By Lemma 2.2 we have $w = w_K$ for some $K \subset \{1, \dots, n-1\}$. $\square$

Definition 2.4. Let $\Delta_j \in \mathbb{C}[(U^+)^e]$ be the top right-hand corner $(n-j) \times (n-j)$ -minor of u in (2.1),

$$(2.3) \quad \Delta_j = \det \begin{pmatrix} x_j & x_{j+1} & & & x_{n-1} \\ x_{j-1} & x_j & \ddots & & \\ & \ddots & \ddots & \ddots & \\ & & \ddots & x_j & x_{j+1} \\ x_{2j-n+1} & & & x_{j-1} & x_j \end{pmatrix},$$

where we set $x_0 = 1$ and $x_k = 0$ for $k < 0$.

We can now give an explicit description of the parts in the decomposition (2.2) of $(U^+)^e$.

Lemma 2.5. For $u \in (U^+)^e$ define $K_u := \{j \in \{1, \dots, n-1\} \mid \Delta_j(u) = 0\}$. Then $u \in B^- w^{K_u} B^-$. In particular,

$$(2.4) \quad (U^+)^e \cap \overline{B^- w^K B^-} = \{u \in (U^+)^e \mid \Delta_j(u) = 0 \text{ all } j \in K\}.$$

Proof. By Lemma 2.3 we have $u \in B^- w^K B^-$ for some $K \subset \{1, \dots, n-1\}$. We need only to show that $K = K_u$. Let $v_1, \dots, v_n$ be the standard basis of $\mathbb{C}^n$ and $\{v_{i_1} \wedge v_{i_2} \wedge \dots \wedge v_{i_d} \mid 1 \leq i_1 < \dots < i_d \leq n\}$ the corresponding basis of $\bigwedge^d \mathbb{C}^n$. Then $\Delta_j(u)$ is the matrix coefficient

$$\Delta_j(u) = (u \cdot (v_{j+1} \wedge \dots \wedge v_n), v_1 \wedge \dots \wedge v_{n-j})$$

of u expressed in this standard basis. Write $u = b_1 w_0 w_K b_2$ for some $b_1 \in B^-$ and $b_2 \in U^-$. Then we have $u \cdot (v_{j+1} \wedge \dots \wedge v_n) = \pm b_1 w_0 w_K \cdot (v_{j+1} \wedge \dots \wedge v_n)$. Therefore $\Delta_j(u) \neq 0$ precisely if $w_0 w_K \cdot (v_{j+1} \wedge \dots \wedge v_n) = \pm v_1 \wedge \dots \wedge v_{n-j}$, or equivalently if

$$(2.5) \quad w_K \cdot (v_{j+1} \wedge \dots \wedge v_n) = \pm v_{j+1} \wedge \dots \wedge v_n.$$

Now consider the maximal parabolic subgroup $W_{\{j\}}$. It can be characterized as the group of permutation matrices preserving the subsets $\{v_1, \dots, v_j\}$ and $\{v_{j+1}, \dots, v_n\}$ in $\mathbb{C}^n$. Therefore the condition (2.5) is equivalent to $w_K \in W_{\{j\}}$, that is, to $j \notin K$. $\square$

Remark 2.6. The preceding lemmas can be generalized to arbitrary reductive linear algebraic groups (where the Δ_j 's are replaced with the corresponding matrix coefficients in the fundamental representations).

3. THE VARIETY $\mathcal{V}_{d,n}$ AND ITS COORDINATE RING

We now go over to studying in detail the one-dimensional components of $(U^+)^e$, corresponding to $K = \{\hat{d}\}$. These are very special in that their elements – and especially their totally positive elements – can be constructed explicitly and in an elementary way. Studying the higher dimensional components involves much deeper results, and this more general case will be treated in the forthcoming paper [17].

Definition 3.1 (The variety $\mathcal{V}_{d,n}$). Define $\mathcal{V}_{d,n}$ to be the closed subvariety of $GL_n(\mathbb{C})$ defined as

$$\mathcal{V}_{d,n} := (U^+)^e \cap \overline{B^- w^{\{\hat{d}\}} B^-},$$

where $w^{\{\hat{d}\}}$ is the Weyl group element defined in 2.3. Let $\mathbb{C}[\mathcal{V}_{d,n}]$ denote the coordinate ring of $\mathcal{V}_{d,n}$ as a reduced affine algebraic variety.

Definition 3.2 (The ring $\Lambda_{d,n}$). For $k = 1, \dots, n$ set $Y_k \in \mathbb{Z}[X_1, \dots, X_d]$ to be the $k \times k$ determinant

$$(3.1) \quad \begin{vmatrix} X_1 & X_2 & \cdots & & \\ & 1 & X_1 & \ddots & \\ & & \ddots & \ddots & \ddots & \vdots \\ & & & \ddots & X_1 & X_2 \\ & & & & 1 & X_1 \end{vmatrix}.$$

In other words, $Y_k = \det(X_{i-j+1})_{i,j=1,\dots,k}$, where $X_0 := 1$ and $X_k := 0$ if $k \notin \{0, \dots, d\}$. We define $\Lambda_{d,n}$ to be the ring given in terms of generators and relations by

$$\Lambda_{d,n} := \mathbb{C}[X_1, \dots, X_d] / (Y_{c+1}, \dots, Y_{n-1}),$$

where $c = n - d$.

Remark 3.3 (The element q and quantum cohomology). Define $q := (-1)^{d+1} Y_n$ in $\Lambda_{d,n}$. Then we may identify

$$\Lambda_{d,n} \cong \mathbb{C}[X_1, \dots, X_d, q] / (Y_{c+1}, \dots, Y_{n-1}, Y_n + (-1)^d q).$$

Therefore the ring $\Lambda_{d,n}$ coincides with $qH^*(Gr_d(n), \mathbb{C})$, the quantum cohomology ring of the Grassmannian of d -planes in $\mathbb{C}^n$, by the presentation found in [19, 20]. See Section 10 for more on this.

By a theorem of Dale Peterson's [16] the quantum cohomology rings of Grassmannians are identified with coordinate rings of the varieties $\mathcal{V}_{d,n}$ (in fact, his statement is a generalization of this to arbitrary type). Using the known presentation of these quantum cohomology rings in type A , Peterson's result for that case may be stated as follows.

Theorem 3.4 (D. Peterson). *There is an isomorphism $\Lambda_{d,n} \xrightarrow{\phi} \mathbb{C}[\mathcal{V}_{d,n}]$ which takes X_j to x_j for $j = 1, \dots, d$.*

Our first aim is to give a direct proof of this theorem. We will not make use of any of the properties of quantum cohomology until Section 11, keeping our treatment completely elementary. So in particular any facts about $\Lambda_{d,n}$ which will be required will be proved by hand, even if they can be deduced from the isomorphism with quantum cohomology.

Lemma 3.5. *The algebra homomorphism $\mathbb{C}[X_1, \dots, X_d] \rightarrow \mathbb{C}[\mathcal{V}_{d,n}]$ defined by $X_j \mapsto x_j$ is surjective and has kernel the radical of the ideal generated by the elements $Y_{c+1}, \dots, Y_{n-1}$ defined in 3.2 above.*

Proof. Let $y_1, \dots, y_n \in \mathbb{C}[(U^+)^e]$ be defined by $y_k := \det(x_{i-j+1})_{i,j=1,\dots,k}$, where $x_0 := 1$ and $x_j := 0$ if $j \notin \{0, \dots, n\}$. It suffices to show that $\mathcal{V}_{d,n}$ is the vanishing set of $x_{d+1}, \dots, x_{n-1}, y_{c+1}, \dots, y_{n-1}$.

By Lemma 2.5, the variety $\mathcal{V}_{d,n}$ consists of the matrices in $(U^+)^e$ for which all the minors Δ_j with $j \neq d$ vanish (see (2.3)).

Consider the minors $\Delta_{n-1}, \Delta_{n-2}, \dots, \Delta_{d+1}$. Note that $\Delta_{n-1} = x_{n-1}$. The vanishing of the minors $\Delta_{n-1}, \dots, \Delta_{d+1}$ implies inductively that the coordinates $x_{d+1}, \dots, x_{n-1}$ vanish. The converse implication is immediate.

Let u be the matrix from (2.1). It is clear that the inverse matrix to u is given by

$$u^{-1} = \begin{pmatrix} 1 & -y_1 & y_2 & \cdots & (-1)^d y_d & \cdots & (-1)^{n-1} y_{n-1} \\ & 1 & -y_1 & & & \ddots & \vdots \\ & & \ddots & \ddots & & & \\ & & & & & & (-1)^d y_d \\ & & & & & & \vdots \\ & & & & 1 & -y_1 & \\ & & & & & & 1 \end{pmatrix}.$$

So $\Delta_1(u) = y_{n-1} = (-1)^{n-1} \Delta_{n-1}(u^{-1})$. Moreover, $\Delta_j(u) = (-1)^{n-1} \Delta_{n-j}(u^{-1})$ for all $j = 1, \dots, n-1$ (see [12] I. (2.9)).

Therefore the conditions $\Delta_1(u) = \dots = \Delta_{d-1}(u) = 0$ are equivalent to $\Delta_{c+1}(u^{-1}) = \Delta_{c+2}(u^{-1}) = \dots = \Delta_{n-1}(u^{-1}) = 0$, and these are equivalent to the conditions $y_{c+1} = \dots = y_{n-1} = 0$. $\square$

Remark 3.6. It follows from the lemma that any element in $\Lambda_{d,n}$ gives rise to a function on $\mathcal{V}_{d,n}$, since we now have a well-defined map $\phi : \Lambda_{d,n} \rightarrow \mathbb{C}[\mathcal{V}_{d,n}]$. We will make free use of this by letting

$$X(u) := \phi(X)(u) \quad \text{for any } X \in \Lambda_{d,n} \text{ and } u \in \mathcal{V}_{d,n}.$$

To fully prove Theorem 3.4, it remains to show that $\Lambda_{d,n}^{\mathbb{C}}$ is reduced. This will be done in Section 5.2.

3.1. Constructing elements in $\mathcal{V}_{d,n}$. We would now like to construct explicitly elements inside $\mathcal{V}_{d,n}$. For $m_1, \dots, m_d \in \mathbb{C}$, let

$$(3.2) \quad \bar{u}_n(m_1, \dots, m_d) := \begin{pmatrix} 1 & m_1 & m_2 & \cdots & m_d & 0 & 0 \\ & 1 & m_1 & \ddots & & \ddots & \\ & & 1 & \ddots & & \ddots & 0 \\ & & & \ddots & & & m_d \\ & & & & \ddots & \ddots & \vdots \\ & & & & & 1 & m_1 & m_2 \\ & & & & & & 1 & m_1 \\ & & & & & & & 1 \end{pmatrix} \in GL_n(\mathbb{C}).$$

Consider the roots $z_1, \dots, z_d \in \mathbb{C}$ of the polynomial

$$z^d - m_1 z^{d-1} + \dots + (-1)^d m_d = 0,$$

counted with multiplicities. So $m_j = E_j(z_1, \dots, z_d)$, where E_j is the j -th elementary symmetric polynomial in $z_1, \dots, z_d$. We define

$$(3.3) \quad u_n(z_1, \dots, z_d) = \bar{u}_n(E_1(z_1, \dots, z_d), \dots, E_d(z_1, \dots, z_d)).$$

There is now a nice description for when a matrix $\bar{u}_n(m_1, \dots, m_d)$ lies in $\mathcal{V}_{d,n}$ in terms of the roots $z_1, \dots, z_d$.

Lemma 3.7. *The matrix $u_n(z_1, \dots, z_d)$ for complex numbers $z_1, \dots, z_d$ lies in $\mathcal{V}_{d,n}$ if and only if either $z_1 = \dots = z_d = 0$ or $z_1, \dots, z_d$ are all distinct and $z_1^n = \dots = z_d^n$. All elements of $\mathcal{V}_{d,n}$ are of this form. Furthermore, the element $q \in \Lambda_{d,n}$ defined in Remark 3.3 evaluates on $u_n(z_1, \dots, z_d) \in \mathcal{V}_{d,n}$ to $q(u_n(z_1, \dots, z_d)) = (-1)^{d+1} z_1^n$.*

The main argument in this lemma could be proved using the presentation of the quantum cohomology ring via the Landau–Ginzburg potential, see [13] Section 8.4, or more originally the work of Gepner [6] for $q = 1$. Below is another proof.

Proof. Let H_k be the k -th complete elementary symmetric polynomial. Then by the Jacobi–Trudi identity, $y_k(u_n(z_1, \dots, z_d)) = H_k(z_1, \dots, z_d)$. So by Lemma 3.5 $\mathcal{V}_{d,n}$ consists of all $u_n(z_1, \dots, z_d)$ for which

$$H_{c+1}(z_1, \dots, z_d) = \dots = H_{n-1}(z_1, \dots, z_d) = 0.$$

Consider the recursion $H_m(x_1, \dots, x_k) = H_m(x_1, \dots, x_{k-1}) + x_k H_{m-1}(x_1, \dots, x_k)$. Therefore

$$(3.4) \quad H_{m-1}(x_1, \dots, x_k) = 0 \implies H_m(x_1, \dots, x_k) = H_m(x_1, \dots, x_{k-1}),$$

and if the x_i are all nonzero, the opposite implication holds as well. Now suppose $H_{c+1}(z_1, \dots, z_d) = \dots = H_{n-1}(z_1, \dots, z_d) = 0$ for $z_1, \dots, z_d \in \mathbb{C}$. Then the vanishing of H_{c+1} implies

$$H_{c+2}(z_1, \dots, z_d) = H_{c+2}(z_1, \dots, z_{d-1}).$$

If $d > 2$ this again equals to zero and it follows that

$$H_{c+3}(z_1, \dots, z_d) = H_{c+3}(z_1, \dots, z_{c-1}) = H_{c+3}(z_1, \dots, z_{d-2}).$$

We repeat this process until we finally get that

$$H_n(z_1, \dots, z_d) = H_n(z_1, \dots, z_{d-1}) = \dots = H_n(z_1) = z_1^n.$$

This implies by symmetry that $z_1^n = z_2^n = \dots = z_d^n = H_n(z_1, \dots, z_d)$. Suppose one of the $z_i \neq 0$. Then the same holds for the others. To prove that the z_i must be distinct, look one step back. We have $0 = H_{n-1}(z_1, \dots, z_d) = H_{n-1}(z_k, z_l)$ for any $k \neq l \in \{1, \dots, d\}$. But $H_{n-1}(z, z) = nz^{n-1}$ is nonzero for $z \neq 0$. Hence $z_k \neq z_l$.

Now suppose we are given d distinct $z_1, \dots, z_d$ with $z_1^n = \dots = z_d^n$. It remains to check that $H_{c+1}, \dots, H_{n-1}$ vanish, or equivalently (since the z_i are nonzero) that $H_n(z_1, \dots, z_d) = \dots = H_n(z_2, z_1) = H_n(z_1) = z_1^n$. We claim that

$$H_{m+n}(z_1, \dots, z_d) = z_1^n H_m(z_1, \dots, z_d).$$

This is obvious if $d = 1$. For $d > 1$ it follows by induction on d using $z_1^n = z_2^n$ and the divided difference identity

$$H_{n+m}(z_1, \dots, z_d) = \frac{H_{n+m+1}(z_1, z_3, \dots, z_d) - H_{n+m+1}(z_2, z_3, \dots, z_d)}{z_1 - z_2}.$$

Now for any $k = 1, \dots, d$ we have

$$\begin{aligned} H_n(z_1, \dots, z_k) &= \frac{H_{n+1}(z_1, z_3, \dots, z_k) - H_{n+1}(z_2, \dots, z_k)}{z_1 - z_2} \\ &= z_1^n \frac{H_1(z_1, z_3, \dots, z_k) - H_1(z_2, \dots, z_k)}{z_1 - z_2} = z_1^n. \end{aligned}$$

This concludes the proof. $\square$

4. ORTHOGONALITY FORMULAS FOR SCHUR POLYNOMIALS AT ROOTS OF UNITY

We have already encountered the elementary symmetric polynomials in d variables $E_1, \dots, E_d$ and the complete homogeneous symmetric polynomials $H_1, H_2, \dots$. At this point the other Schur polynomials also enter the picture. Denote by S_λ the Schur polynomial in d variables corresponding to the partition λ of at most d parts. We will write such a partition λ as a d -tuple of integers $\lambda = (\lambda_1, \dots, \lambda_d)$ with $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_d \geq 0$. As a visual aid partitions are often represented by their Young diagrams, see [12].

There are many definitions of the Schur polynomials. For instance, the Schur polynomials are the characters of the polynomial representations of GL_d (symmetric polynomials in the eigenvalues). We recall also that the Schur polynomials may be obtained from the elementary and complete homogeneous symmetric polynomials by the Jacobi–Trudi and dual Jacobi–Trudi identities: Let $\lambda^t = (\lambda_1^t, \dots, \lambda_c^t)$ be the conjugate partition to λ obtained by exchanging rows and columns in the Young diagram, then

$$S_\lambda = \begin{vmatrix} H_{\lambda_1} & H_{\lambda_1+1} & \cdots & H_{\lambda_1+d-1} \\ H_{\lambda_2-1} & H_{\lambda_2} & \cdots & H_{\lambda_2+d-2} \\ & & \ddots & \\ H_{\lambda_d-d+1} & \cdots & \cdots & H_{\lambda_d} \end{vmatrix} = \begin{vmatrix} E_{\lambda_1^t} & E_{\lambda_1^t+1} & \cdots & E_{\lambda_1^t+c-1} \\ E_{\lambda_2^t-1} & E_{\lambda_2^t} & \cdots & E_{\lambda_2^t+c-2} \\ & & \ddots & \\ E_{\lambda_c^t+c-1} & \cdots & \cdots & E_{\lambda_c^t} \end{vmatrix},$$

where $E_k = H_k = 0$ for $k < 0$. The Schur polynomial S_λ is homogeneous of degree $|\lambda| := \lambda_1 + \dots + \lambda_d$, the size of the partition. Also, the Schur polynomials form a basis of the ring of symmetric polynomials in d variables. For further background on Schur polynomials we refer to [12].

Definition 4.1 (partitions inside a box and PD). Let $\text{Sh}(d, c)$ denote the set of partitions whose Young diagram or shape fits into a $d \times c$ box. In other words $\text{Sh}(d, c)$ is the set of partitions $\lambda = (\lambda_1, \dots, \lambda_d)$ of length at most d such that $\lambda_1 \leq c$. We use the shorthand notation (m^k) for the partition $(m, \dots, m, 0, \dots, 0)$ with m occurring k times. So for example (c^d) is the longest partition in $\text{Sh}(d, c)$.

For a partition $\lambda \in \text{Sh}(d, c)$ define

$$\text{PD}(\lambda) := (c - \lambda_d, c - \lambda_{d-1}, \dots, c - \lambda_1).$$

The notation PD stands for ‘Poincaré duality’ (see Section 10).

Definition 4.2 (The set $\mathcal{I}_{d,n}$). We fix the primitive n -th root of unity $\zeta = \exp(\frac{2\pi i}{n})$. Let $\zeta^I := (\zeta^{i_1}, \dots, \zeta^{i_d})$ be an unordered d -tuple of distinct n -th roots of $(-1)^{d+1}$. Then $I = (i_1, \dots, i_d)$ may be chosen uniquely such that $-\frac{d-1}{2} \leq i_1 < i_2 < \dots < i_d \leq n - \frac{d+1}{2}$ and the i_k ’s are all integers (respectively half-integers) if d is odd (even). Denote the set of all such d -tuples I by $\mathcal{I}_{d,n}$.

Let $c = n - d$. If $I = (i_1, \dots, i_d) \in \mathcal{I}_{d,n}$, then denote by $\hat{I} = (\hat{i}_1, \dots, \hat{i}_c)$ the unique c -tuple, $-\frac{d-1}{2} \leq \hat{i}_1 < \hat{i}_2 < \dots < \hat{i}_c \leq n - \frac{d+1}{2}$, such that $\zeta^{i_1}, \dots, \zeta^{i_d}, \zeta^{\hat{i}_1}, \dots, \zeta^{\hat{i}_c}$ enumerates all n -th roots of $(-1)^{d+1}$.

There is a bijection

$$(4.1) \quad (\cdot)^t : \mathcal{I}_{d,n} \rightarrow \mathcal{I}_{c,n}$$

which takes $I = (i_1, \dots, i_d)$ to $I^t := (\frac{n}{2} - \hat{i}_c, \dots, \frac{n}{2} - \hat{i}_1)$.

We note that this bijection corresponds exactly to transposition of shapes, $(\cdot)^t : \text{Sh}(d, c) \rightarrow \text{Sh}(c, d)$, after the identification

$$(4.2) \quad \begin{array}{ccc} \text{Sh}(d, c) & \leftrightarrow & \mathcal{I}_{d,n} \\ \lambda = (\lambda_1, \dots, \lambda_d) & \mapsto & I_\lambda = (\frac{d+1}{2} + \lambda_d - d, \dots, \frac{d+1}{2} + \lambda_1 - 1). \end{array}$$

We have $I_{\lambda^t} = (I_\lambda)^t$. Furthermore if $\|I\| := \sum_{k=1}^d i_k$ then $\|I_\lambda\| = |\lambda|$.

We will show the following identities for Schur polynomials.

Proposition 4.3 (Orthogonality formulas). *Let $I, J \in \mathcal{I}_{d,n}$ and $z_1, \dots, z_d, t \in \mathbb{C}$. Then*

$$\begin{aligned} (1) \quad & \sum_{\lambda \in \text{Sh}(d, c)} S_\lambda(z_1, \dots, z_d) S_{\text{PD}(\lambda)}(\zeta^J) = S_{(c^d)}(\zeta^J) \prod_{k=1}^d \prod_{l=1}^c (1 - z_k \zeta^{-\hat{j}_l}), \\ (2) \quad & \sum_{\lambda \in \text{Sh}(d, c)} S_\lambda(t \zeta^I) S_{\text{PD}(\lambda)}(t \zeta^J) = \delta_{I, J} \frac{n^d S_{(c^d)}(t \zeta^I)}{|\text{Vand}(\zeta^I)|^2}, \\ (3) \quad & \sum_{\lambda \in \text{Sh}(d, c)} S_\lambda(\zeta^I) \overline{S_\lambda(\zeta^J)} = \delta_{I, J} \frac{n^d}{|\text{Vand}(\zeta^I)|^2}. \end{aligned}$$

where $|\text{Vand}(\zeta^I)|$ stands for the absolute value of the Vandermonde determinant $\prod_{k < j} (\zeta^{i_k} - \zeta^{i_j})$ and the bar in equation (3) stands for complex conjugation.

Lemma 4.4. *If $\lambda \in \text{Sh}(d, c)$ and $I \in \mathcal{I}_{d,n}$ then we have the following equality of values of Schur polynomials,*

$$(4.3) \quad S_{\lambda^t}(\zeta^{I^t}) = \frac{S_{\text{PD}(\lambda)}(\zeta^I)}{S_{(c^d)}(\zeta^I)} = \overline{S_\lambda(\zeta^I)}.$$

Proof. To begin with, recall that $S_{(c^d)}$ is just the character of the c -th power of the determinant representation of GL_d . The right hand side equality follows from $\overline{S_\lambda(\zeta^I)} = S_\lambda(\zeta^{-I})$ and the general formula

$$S_\lambda(z_1^{-1}, \dots, z_d^{-1}) = \frac{S_{\text{PD}(\lambda)}(z_1, \dots, z_d)}{S_{(c^d)}(z_1, \dots, z_d)}$$

for the character of the dual representation of GL_d .

We now prove that $S_{\lambda^t}(\zeta^{\frac{n}{2} - \hat{I}}) = S_\lambda(\zeta^{-I})$. Let $1 \leq k \leq d$ and suppose $\lambda \in \text{Sh}(d, c)$ is the partition $(1^k) := (1, \dots, 1, 0, \dots, 0)$ with 1 appearing k times. So $S_\lambda = E_k$, the k -th elementary symmetric polynomial, and $S_{\lambda^t} = H_k$, the k -th

complete homogeneous symmetric polynomial. Then we have

$$\begin{aligned} p(t) &:= \prod_{j=1}^d (1 + t\zeta^{-i_j}) = 1 + E_1(\zeta^{-I})t + E_2(\zeta^{-I})t^2 + \dots + E_d(\zeta^{-I})t^d, \\ r(t) &:= \prod_{j=1}^d \frac{1}{(1 - t\zeta^{\frac{n}{2} - i_j})} = 1 + H_1(\zeta^{\frac{n}{2} - \hat{I}})t + H_2(\zeta^{\frac{n}{2} - \hat{I}})t^2 + \dots + H_d(\zeta^{\frac{n}{2} - \hat{I}})t^d + 0 + \\ &\quad + 0 + H_n(\zeta^{\frac{n}{2} - \hat{I}})t^n + \text{higher order terms.} \end{aligned}$$

And since

$$r(t)^{-1}p(t) = \prod_{k=1}^n (1 + t\zeta^{-\frac{d+1}{2} + k}) = 1 + (-1)^{d+1}t^n$$

we get $r(t)(1 + (-1)^{d+1}t^n) = p(t)$. Comparing the first d coefficients on either side we see that $E_k(\zeta^{-I}) = H_k(\zeta^{\frac{n}{2} - \hat{I}})$. So the required identity is proved for $\lambda = (1^k)$, where $k = 1, \dots, d$. For all other partitions in $\text{Sh}(d, c)$ the formula follows from this special case using the Jacobi–Trudi identity. $\square$

Proof of Proposition 4.3. Recall the classical identity due to Littlewood,

$$(4.4) \quad \sum_{\lambda \in \text{Sh}(d, c)} S_\lambda(z_1, \dots, z_d) S_{\lambda^t}(w_1, \dots, w_c) = \prod_{i=1}^d \prod_{j=1}^c (1 + z_i w_j),$$

which is a consequence of the usual orthogonality of Schur functions (see Macdonald [12] I.(4.3'), or Fulton [4] Appendix A). Now replace $(w_1, \dots, w_c)$ by ζ^{J^t} in (4.4) and apply Lemma 4.4 to get the orthogonality formula (1). Formulas (2) and (3) follow by furthermore replacing $(z_1, \dots, z_d)$ by ζ^I and checking that

$$\prod_{k=1}^d \prod_{l=1}^c (1 - \zeta^{i_k} \zeta^{-\hat{j}_l}) = \begin{cases} 0 & \text{if } I \neq J \\ \frac{n^d}{|\text{Vand}(\zeta^I)|^2} & \text{if } I = J. \end{cases}$$

This is immediate in the case $I \neq J$. To verify the identity for $I = J$ compute

$$|\text{Vand}(\zeta^I)|^2 \prod_{k=1}^d \prod_{l=1}^c (1 - \zeta^{i_k} \zeta^{-\hat{i}_l}) = \prod_{m, r \in \{1, \dots, d\}, m \neq r} (1 - \zeta^{i_m} \zeta^{-i_r}) \prod_{k=1}^d \prod_{l=1}^c (1 - \zeta^{i_k} \zeta^{-\hat{i}_l}),$$

which is the same as

$$(4.5) \quad \prod_{k=1}^d \left[\frac{\prod_{r=1}^n (1 - z\zeta^{i_k} \zeta^{r - \frac{d+1}{2}})}{1 - z} \right]_{z=1}.$$

Since this polynomial in z simplifies to

$$p(z) = \frac{\prod_{l=1}^n (1 - z\zeta^{i_k} \zeta^{-\frac{d+1}{2} + l})}{1 - z} = \frac{1 - z^n}{1 - z} = 1 + z + \dots + z^{n-1},$$

we have $p(1) = n$ and (4.5) equals to n^d . $\square$

Remark 4.5. Passing from ‘column orthogonality’ to ‘row orthogonality’ in Proposition 4.3 we also get the formulas

$$(4.6) \quad \frac{1}{n^d} \sum_{I \in \mathcal{I}_{d,n}} S_\lambda(\zeta^I) \overline{S_\mu(\zeta^I)} |\text{Vand}(\zeta^I)|^2 = \delta_{\lambda,\mu}$$

and

$$(4.7) \quad \frac{1}{n^d} \sum_{I \in \mathcal{I}_{d,n}} S_\lambda(\zeta^I) S_{\text{PD}(\mu)}(\zeta^I) \frac{|\text{Vand}(\zeta^I)|^2}{S_{(c^d)}(\zeta^I)} = \delta_{\lambda,\mu},$$

for any $\lambda, \mu \in \text{Sh}(d, c)$. The first one, (4.6), looks like it should have an explanation on terms of representation theory of GL_d . The second one is related to the analogue of Poincaré duality in the quantum cohomology ring, see Section 10.5.

5. APPLICATIONS TO THE RING $\Lambda_{d,n}$

In this section we will finish the proof of Theorem 3.4, that $\Lambda_{d,n}$ is the coordinate ring of $\mathcal{V}_{d,n}$. For a partition $\lambda \in \text{Sh}(d, c)$, let $s_\lambda \in \Lambda_{d,n}$ be defined by

$$(5.1) \quad s_\lambda = \begin{vmatrix} X_{\lambda_1^t} & X_{\lambda_1^t+1} & \cdots & X_{\lambda_1^t+c-1} \\ X_{\lambda_2^t-1} & X_{\lambda_2^t} & \cdots & X_{\lambda_2^t+c-2} \\ & & \ddots & \\ X_{\lambda_c^t+c-1} & \cdots & \cdots & X_{\lambda_c^t} \end{vmatrix}.$$

Note that this looks like the dual Jacobi–Trudi formula. Therefore these elements evaluate on a point $u = u_n(z_1, \dots, z_d)$ in $\mathcal{V}_{d,n}$ to $s_\lambda(u) = S_\lambda(z_1, \dots, z_d)$.

5.1. The quantum Pieri rule. We will need to know how these s_λ behave under multiplication by the generators X_k of $\Lambda_{d,n}$. This is explained by the (dual) quantum Pieri rule stated below. The quantum Pieri rule was first found by Bertram [2] who proved it for the ring $qH^*(Gr_d(n))$ using some geometry. But later Bertram, Ciocan–Fontanine, and Fulton [3] also gave an algebraic and combinatorial proof inside the ring $\Lambda_{d,n}$.

Theorem 5.1 (dual quantum Pieri rule, [3] Proposition 4.2). *Let $\lambda \in \text{Sh}(d, c)$ and $k \in \{1, \dots, d\}$, then the following formula holds in $\Lambda_{d,n}$,*

$$X_k s_\lambda = \sum_{\nu} s_\nu + q \sum_{\mu} s_\mu,$$

where first sum is over all $\nu \in \text{Sh}(d, c)$ such that $|\nu| = |\lambda| + k$ and $\nu_1^t \geq \lambda_1^t \geq \nu_2^t \geq \dots \geq \nu_c^t \geq \lambda_c^t \geq 0$. The second sum is over all μ such that $|\mu| = |\lambda| + k - n$ and $\lambda_1^t - 1 \geq \mu_1^t \geq \lambda_2^t - 1 \geq \mu_2^t \geq \dots \geq \lambda_d^t - 1 \geq \nu_1^t \geq 0$.

Such a formula for multiplication at $q = 1$ also shows up in the earlier work [6] of Gepner coming from physics. And Gepner notes that this specialized ring is semi-simple. The final step of our proof of Theorem 3.4 below uses a similar insight to Gepner’s.

5.2. Conclusion of the proof of Theorem 3.4. By Lemma 3.5 it remains only to prove that the homomorphism $\phi : \Lambda_{d,n}^{\mathbb{C}} \rightarrow \mathbb{C}[\mathcal{V}_{d,n}]$ is injective. In other words, let $p \in \Lambda_{d,n}^{\mathbb{C}}$, and suppose the image of p in $\mathbb{C}[\mathcal{V}_{d,n}]$, again denoted by p , is zero. Then we must show that $p = 0$ in $\Lambda_{d,n}^{\mathbb{C}}$.

At the outset, p is some polynomial in the X_k 's. By repeated application of the dual quantum Pieri rule we can write p as a linear combination $p = \sum_{\lambda \in \text{Sh}(d,c)} p_{\lambda} s_{\lambda}$, where $p_{\lambda} \in \mathbb{C}[q]$. Then because p vanishes as function on $\mathcal{V}_{d,n}$ we get that

$$(5.2) \quad 0 = p(u_n(t\zeta^I)) = \sum_{\lambda} p_{\lambda}(t^n) S_{\lambda}(t\zeta^I),$$

for any element $u_n(t\zeta^I) \in \mathcal{V}_{d,n}$. Now for any fixed $t \in \mathbb{C}$ consider the vector $(p_{\lambda}(t^n))_{\lambda} \in \mathbb{C}^{\text{Sh}(d,c)}$. By (5.2) this vector lies in the orthogonal complement to the span $\langle (S_{\lambda}(t\zeta^I))_{\lambda} \rangle_{I \in \mathcal{I}_{d,n}}$ (with respect to the standard inner product on $\mathbb{C}^{\text{Sh}(d,c)}$).

But the vectors $(S_{\lambda}(t\zeta^I))_{\lambda}$ for varying $I \in \mathcal{I}_{d,n}$ are all linearly independent, by the orthogonality formula in Proposition 4.3 (2), and therefore span $\mathbb{C}^{\text{Sh}(d,c)}$. So it follows that the vector $(p_{\lambda}(t^n))_{\lambda} = 0$. Since this holds for all $t \in \mathbb{C}$ we have that all of the $p_{\lambda} = 0$ in $\mathbb{C}[q]$. Hence $p = 0$ as required. $\square$

Corollary 5.2 (Schubert basis). *The elements $\{q^k s_{\lambda} \mid k \in \mathbb{Z}_{\geq 0}, \lambda \in \text{Sh}(d,c)\}$ form a $\mathbb{Z}$ -basis of $\Lambda_{d,n}^{\mathbb{Z}} = \mathbb{Z}[X_1, \dots, X_d]/(Y_{c+1}, \dots, Y_{n-1})$.*

Proof. By repeated application of the quantum Pieri rule any element in $\Lambda_{d,n}^{\mathbb{Z}}$ is expressible as a $\mathbb{Z}$ -linear combination of elements $q^k s_{\lambda}$. These are all linearly independent by exactly the same argument as in the proof of Theorem 3.4 above. $\square$

6. STRUCTURE CONSTANTS OF $\Lambda_{d,n}$

In this section we give an elementary proof of a formula for the structure constants of $\Lambda_{d,n}$. The formula (6.1) below was first noted by Bertram [2] to hold in the isomorphic ring $qH^*(Gr_d(n))$ as a consequence of his quantum Giambelli formula (Theorem 10.2) and the Vafa–Intriligator formula for Gromov–Witten invariants. The mathematical proof of the Vafa–Intriligator formula by Siebert and Tian [19] depends on higher-dimensional residues and the geometric definition of the structure constants. We give a simple proof for $\Lambda_{d,n}$ below which instead uses Theorem 3.4 and the orthogonality formula from Proposition 4.3.

Proposition 6.1. *Let $\lambda_1, \dots, \lambda_N \in \text{Sh}(d,c)$, $I \in \mathcal{I}_{d,n}$ and $t \in \mathbb{C}^*$. Let P be an arbitrary homogeneous symmetric polynomial in d variables.*

1. *We have the following formula for $P(t\zeta^I)$ in terms of Schur polynomials,*

$$P(t\zeta^I) = \sum_{\nu \in \text{Sh}(d,c)} m_{\nu}^P(t) S_{\nu}(t\zeta^I)$$

$$\text{where} \quad m_{\nu}^P(t) = \frac{1}{n^d} \sum_{J \in \mathcal{I}_{d,n}} P(t\zeta^J) S_{\nu}(t^{-1}\zeta^{-J}) |\text{Vand}(\zeta^J)|^2.$$

2. *For any homogeneous symmetric polynomial m ,*

$$\sum_{J \in \mathcal{I}_{d,n}} m(\zeta^J) = 0 \quad \text{unless} \quad \deg(m) \equiv 0 \pmod{n}.$$

In particular, $m_{\nu}^P = 0$ unless $|\nu| \equiv \deg(P) \pmod{n}$.

3. Let p be a (non-zero) homogeneous element in $\Lambda_{d,n}$. Then the coefficients $p_{\lambda,k}$ of

$$p = \sum_{\lambda \in \text{Sh}(d,c), k \in \mathbb{Z}_{\geq 0}} p_{\lambda,k} q^k s_{\lambda}$$

written out in the basis $\{q^k s_{\lambda} \mid k \in \mathbb{Z}_{\geq 0}, \lambda \in \text{Sh}(d,c)\}$ of $\Lambda_{d,n}$ are given by the formula

$$p_{\lambda,k} = \begin{cases} \frac{1}{n^d} \sum_{J \in \mathcal{I}_{d,n}} p(u(\zeta^J)) \overline{S_{\lambda}(\zeta^J)} |\text{Vand}(\zeta^J)|^2 & \text{if } kn + |\lambda| = \deg(p), \\ 0 & \text{otherwise.} \end{cases}$$

Corollary 6.2 (Bertram–Vafa–Intriligator Formula). *The structure constants for multiplication in the basis $\{q^k s_{\lambda} \mid \lambda \in \text{Sh}(d,c), k \in \mathbb{N}\}$ of $\Lambda_{d,n}$,*

$$s_{\lambda} s_{\mu} = \sum_{k \in \mathbb{N}, \lambda_0 \in \text{Sh}(d,c)} \langle s_{\lambda}, s_{\mu}, s_{\nu} \rangle_k q^k s_{\text{PD}(\nu)},$$

are computed by

$$(6.1) \quad \langle s_{\lambda}, s_{\mu}, s_{\nu} \rangle_k = \frac{1}{n^d} \sum_{J \in \mathcal{I}_{d,n}} S_{\lambda}(\zeta^J) S_{\mu}(\zeta^J) S_{\nu}(\zeta^J) \frac{|\text{Vand}(\zeta^J)|^2}{S_{(cd)}(\zeta^J)}$$

whenever $cd + kn = |\lambda| + |\mu| + |\nu|$, and otherwise by $\langle s_{\lambda}, s_{\mu}, s_{\nu} \rangle_k = 0$.

Proof of the Corollary. The formula (6.1) is an immediate consequence of Proposition 6.1(3) and Lemma 4.4. $\square$

Proof of Proposition 6.1. For $J \in \mathcal{I}_{d,n}$ define the symmetric polynomial $S_J := \sum_{\nu \in \text{Sh}(d,c)} S_{\nu}(\zeta^{-J}) S_{\nu}$. By Theorem 4.3(3) we have $S_J(\zeta^I) = \delta_{I,J} \frac{n^d}{|\text{Vand}(\zeta^J)|^2}$. Therefore,

$$\begin{aligned} P(t\zeta^I) &= \frac{1}{n^d} \sum_{J \in \mathcal{I}_{d,n}} P(t\zeta^J) |\text{Vand}(\zeta^J)|^2 S_J(\zeta^I) = \\ &= \frac{1}{n^d} \sum_{\nu \in \text{Sh}(d,c)} \sum_{J \in \mathcal{I}_{d,n}} P(t\zeta^J) |\text{Vand}(\zeta^J)|^2 S_{\nu}(t^{-1}\zeta^{-J}) S_{\nu}(t\zeta^I). \end{aligned}$$

So (1) is proved.

For (2) notice that $M(t) := \sum_{J \in \mathcal{I}_{d,n}} m(t\zeta^J)$ satisfies $M(t) = M(\zeta t)$. Therefore M is a polynomial in t^n , and unless n divides $\deg(m)$ we must have $M = 0$. Since $(\deg(P) - |\nu|)$ is the degree of m_{ν}^P , the rest of (2) also follows.

To prove any identity in $\Lambda_{d,n}$ it suffices, by Theorem 3.4, to check it ‘on points’, that is, evaluated at the elements of $\mathcal{V}_{d,n}$. Therefore we can deduce (3) directly from (1) and (2). $\square$

Remark 6.3. The ring $\Lambda_{d,n}$ specializes to $\Lambda_{d,n}^{(q=0)} = \mathbb{C}[x_1, \dots, x_d]/(y_{c+1}, \dots, y_n)$, the usual cohomology ring of the Grassmannian, at $q = 0$. The (images of the) s_{λ} with $\lambda \in \text{Sh}(d,c)$ now form a basis of this ring, and the structure constants for this basis are the well-known Littlewood–Richardson coefficients. Therefore the formula (6.1) for structure constants in $\Lambda_{d,n}$ gives when $k = 0$ a formula for Littlewood–Richardson coefficients.

7. TOTAL POSITIVITY (A QUICK REVIEW)

Before proceeding in the next section to determining the totally positive part of $\mathcal{V}_{d,n}$ we will give some background and motivation for doing this.

Total positivity, introduced by Schoenberg in the 1930's is the study of invertible matrices with exclusively nonnegative real minors (and nonnegative entries in particular). We will strictly speaking call such elements of GL_n totally nonnegative, and, if all minors are in fact strictly positive, then totally positive. The definition of total positivity is basis-dependent (or more precisely dependent on a 'pinning' of GL_n , see [10]), but totally positive matrices also have nice intrinsic properties. For example, Gantmacher and Krein showed that all totally positive matrices are regular semisimple with distinct positive real eigenvalues.

The totally positive matrices form an open semialgebraic subset of the real points $GL_n(\mathbb{R})$ which we denote by $GL_n(\mathbb{R}_{>0})$ in analogy with the standard notation $GL_n(\mathbb{K})$ for the $\mathbb{K}$ -valued points of GL_n when $\mathbb{K}$ is a field. We also write $GL_n(\mathbb{R}_{\geq 0})$ for the totally nonnegative part of GL_n .

In the early 1990's Lusztig [10] took the theory of total positivity a big step further and generalized it to all reductive algebraic groups, at the same time relating it (in the simply-laced case) to canonical bases and their deep positivity properties. Let us look in more detail at the special case of the subgroup U^+ , to give a taste of what the totally positive part looks like and how it is related to the canonical basis. We use the notations introduced in 2.1.

Let $U^+(\mathbb{R}) = U^+ \cap GL_n(\mathbb{R})$. We will consider $GL_n(\mathbb{R})$ and any subsets as endowed with the usual Hausdorff topology. Inside $U^+(\mathbb{R})$ we define $U^+(\mathbb{R}_{\geq 0})$ to be the semigroup of all totally nonnegative matrices in U^+ . This is a semigroup since the totally nonnegative matrices can be characterized as acting by nonnegative matrices in all exterior powers of the standard representation of GL_n (with their standard bases).

In the 1950's A. Whitney proved that $U^+(\mathbb{R}_{\geq 0})$ is the semigroup generated by the elements $\exp(te_i)$ for $t \in \mathbb{R}_{\geq 0}$. The following more precise description of $U^+(\mathbb{R}_{\geq 0})$ was given by Lusztig in [10]. For $w \in W$ let $U_w^+ = U^+ \cap B^- w B^-$. Then we define $U_w^+(\mathbb{R}_{\geq 0})$ to be the set of totally nonnegative matrices in U_w^+ . For w equal to the simple reflection s_i it is easy to see that

$$\begin{aligned} \mathbb{R}_{>0} &\xrightarrow{\sim} U_{s_i}^+(\mathbb{R}_{>0}), \\ t &\mapsto x_i(t) := \exp(te_i), \end{aligned}$$

and $U_{s_i}^+(\mathbb{R}_{>0})$ is a connected component of $U_{s_i}^+(\mathbb{R}) \cong \mathbb{R}^*$. Hence the notation $U_{s_i}^+(\mathbb{R}_{>0})$ rather than $U_{s_i}^+(\mathbb{R}_{\geq 0})$.

Moreover, for general w taking a reduced expression $w = s_{i_1} \dots s_{i_k}$ and multiplying together the corresponding maps gives a (semi-algebraic) isomorphism

$$(7.1) \quad \begin{aligned} (\mathbb{R}_{>0})^k &\xrightarrow{\sim} U_w^+(\mathbb{R}_{>0}), \\ (t_1, \dots, t_k) &\mapsto x_{i_1}(t_1) \cdot \dots \cdot x_{i_k}(t_k), \end{aligned}$$

see [10], Proposition 2.7.

The largest of these real semi-algebraic cells, $U_{w_0}^+(\mathbb{R}_{>0})$, which corresponds to the longest element in W , is also called the totally positive part of U^+ and denoted by $U^+(\mathbb{R}_{>0})$. It is open in $U^+(\mathbb{R})$ and its closure is the set of all totally nonnegative matrices in U^+ , denoted by $U^+(\mathbb{R}_{\geq 0})$.

Now let us describe the connection with canonical bases. Consider the coordinate ring $\mathbb{C}[U^+]$ as the graded dual to the enveloping algebra $\mathcal{U}^+$ of the Lie algebra $\text{Lie}(U^+)$. Then $\mathbb{C}[U^+]$ has a basis $\mathbb{B}$ given by the dual canonical basis (obtained from the canonical basis of the dual quantum enveloping algebra after specializing the quantum parameter to 1). In fact it also has a $\mathbb{Z}$ -form which is spanned by this canonical basis. And the products of basis vectors are nonnegative linear combinations of basis vectors: the structure constants are nonnegative integers. This last point is very deep as it follows from Lusztig's geometric construction of the coproduct in the quantized universal enveloping algebra [9]. And this positivity property enters crucially into the proof of the following theorem.

Theorem 7.1 (Lusztig [10], see also [11] 3.13). *Let $u \in U^+$. Then*

$$\begin{aligned} u \in U^+(\mathbb{R}_{>0}) &\iff b(u) \in \mathbb{R}_{>0} \text{ for all } b \in \mathbb{B}, \\ \text{and } u \in U^+(\mathbb{R}_{\geq 0}) &\iff b(u) \in \mathbb{R}_{\geq 0} \text{ for all } b \in \mathbb{B}. \end{aligned}$$

We will see in the next section exactly similar results for the variety $\mathcal{V}_{d,n}$, where the dual canonical basis is replaced by the Schubert basis.

8. THE TOTALLY POSITIVE PART OF $\mathcal{V}_{d,n}$

We begin with some general observations about $\mathcal{V}_{d,n}$. Firstly $\mathcal{V}_{d,n}$ is a 1-dimensional affine variety with a $\mathbb{C}^*$ -action, given explicitly by


$$t \cdot \bar{u}_n(m_1, \dots, m_d) = \bar{u}_n(tm_1, t^2m_2, \dots, t^dm_d)$$

(see (3.2)). We identify $\Lambda_{d,n}$ with $\mathbb{C}[\mathcal{V}_{d,n}]$ by the isomorphism of Theorem 3.4. Consider the morphism therefore defined by $q \in \Lambda_{d,n}$,

$$q : \mathcal{V}_{d,n} \rightarrow \mathbb{C}.$$

This map is $\mathbb{C}^*$ -equivariant for the $\mathbb{C}^*$ -action on $\mathbb{C}$ given by $t \cdot z = t^n z$, for $t \in \mathbb{C}^*$, $z \in \mathbb{C}$. And the fiber over 1 is precisely $\{u_n(\zeta^I) \mid I \in \mathcal{I}_{d,n}\}$. So q is an $\binom{n}{d}$ -fold covering over $\mathbb{C}^*$, ramified at 0. Also, two elements $u_n(\zeta^I)$ and $u_n(\zeta^J)$ in the fiber of q over 1 lie in the same $\mathbb{C}^*$ -orbit if and only if $\zeta^J = \zeta^k \zeta^I$ for some $k = 1, \dots, n$. Therefore $\mathcal{V}_{d,n}$ consists of precisely $\frac{1}{n} \binom{n}{d}$ regular $\mathbb{C}^*$ -orbits, and one trivial one (given by the identity matrix).

Consider now the real form $\Lambda_{d,n}^{\mathbb{R}} = \mathbb{R}[X_1, \dots, X_d] / (Y_{c+1}, \dots, Y_{n-1}) = \Lambda_{d,n}^{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{R}$, and the corresponding set of real points $\mathcal{V}_{d,n}(\mathbb{R}) = \mathcal{V}_{d,n} \cap GL_n(\mathbb{R})$ of $\mathcal{V}_{d,n}$. If $d' := \lfloor \frac{d}{2} \rfloor$ is the greatest integer part of $\frac{d}{2}$, then there are precisely $\binom{n}{d'}$ real points in the fiber of q over 1. So $\mathcal{V}_{d,n}(\mathbb{R})$ is made up of $\binom{n}{d'}$ regular $\mathbb{R}^*$ -orbits and the one trivial one, and $q : \mathcal{V}_{d,n}(\mathbb{R}) \rightarrow \mathbb{R}$ simply looks as indicated in the figure below.



We will identify exactly one of the branches over $\mathbb{R}_{>0}$ as the totally positive part of $\mathcal{V}_{d,n}$. Let us give the precise definitions first.

Definition 8.1. Let $\mathcal{V}_{d,n}(\mathbb{R}_{\geq 0}) := \mathcal{V}_{d,n}(\mathbb{R}) \cap U^+(\mathbb{R}_{\geq 0})$, the set of totally nonnegative matrices in $\mathcal{V}_{d,n}$. Recall that by definition

$$\mathcal{V}_{d,n} = (U^+)^e \cap \overline{(B^- w^{\{d\}} B^-)}.$$

Therefore in analogy with the case of U^+ described in the previous section we define $\mathcal{V}_{d,n}(\mathbb{R}_{>0}) := \mathcal{V}_{d,n}(\mathbb{R}_{\geq 0}) \cap (B^- w^{\{d\}} B^-)$. The remaining Bruhat decomposition of $\mathcal{V}_{d,n}(\mathbb{R}_{\geq 0})$ is very simple since, by Lemma 2.3 or the above figure, we have

$$\mathcal{V}_{d,n}(\mathbb{R}_{\geq 0}) = \mathcal{V}_{d,n}(\mathbb{R}_{>0}) \sqcup \{id\}.$$

Definition 8.2. Let $I_0 = (-\frac{d-1}{2}, -\frac{d-1}{2} + 1, \dots, \frac{d-1}{2}) \in \mathcal{I}_{d,n}$. So ζ^{I_0} is in a sense the ‘positive-most’ d -tuple of roots of $(-1)^{d+1}$. Then we define a one-parameter family of matrices in $\mathcal{V}_{d,n}(\mathbb{R})$ by

$$(8.1) \quad u_{>0} : \mathbb{R} \rightarrow \mathcal{V}_{d,n}(\mathbb{R}), \quad \text{where} \quad u_{>0}(t) := u_n(t\zeta^{I_0}).$$

Note that $u_n(t\zeta^{I_0})$ does indeed lie in $\mathcal{V}_{d,n}(\mathbb{R})$ since the d -tuple ζ^{I_0} is (up to permutation) invariant under complex conjugation.

8.1. Let us determine the matrix entries $E_1(t\zeta^{I_0}), \dots, E_d(t\zeta^{I_0})$ of $u_n(t\zeta^{I_0})$ explicitly. We use a variant of the classical q -binomial theorem of Gauss, with ζ playing the role of q (c.f. [12] I.2 Example 3). For $m, k \in \mathbb{N}$, let

$$[m]_\zeta := \frac{\zeta^{\frac{m}{2}} - \zeta^{-\frac{m}{2}}}{\zeta^{\frac{1}{2}} - \zeta^{-\frac{1}{2}}} \quad \text{and} \quad \begin{bmatrix} m \\ k \end{bmatrix}_\zeta := \frac{[m]_\zeta [m-1]_\zeta \dots [m-k+1]_\zeta}{[1]_\zeta [2]_\zeta \dots [k]_\zeta}.$$

Then we have the following version of the binomial formula,

$$\begin{aligned} \left(1 + \zeta^{-\frac{d-1}{2}} t\right) \left(1 + \zeta^{-\frac{d-1}{2}+1} t\right) \dots \left(1 + \zeta^{\frac{d-1}{2}} t\right) = \\ 1 + \begin{bmatrix} d \\ 1 \end{bmatrix}_\zeta t + \begin{bmatrix} d \\ 2 \end{bmatrix}_\zeta t^2 + \dots + \begin{bmatrix} d \\ d-1 \end{bmatrix}_\zeta t^{d-1} + t^d. \end{aligned}$$

In other words the elementary symmetric polynomials evaluated on $t\zeta^{I_0}$ give

$$E_j \left(t\zeta^{-\frac{d-1}{2}}, t\zeta^{-\frac{d-1}{2}+1}, \dots, t\zeta^{\frac{d-1}{2}} \right) = t^j \begin{bmatrix} d \\ j \end{bmatrix}_\zeta.$$

Since $\zeta = \exp(\frac{2\pi i}{n})$ we have that $\frac{\zeta^{\frac{m}{2}} - \zeta^{-\frac{m}{2}}}{\zeta^{\frac{1}{2}} - \zeta^{-\frac{1}{2}}} = \frac{\sin(\frac{m\pi}{n})}{\sin(\frac{\pi}{n})}$. Therefore

$$E_j(t\zeta^{I_0}) = t^j \frac{\sin(\frac{d\pi}{n}) \sin((d-1)\frac{\pi}{n}) \dots \sin((d-j+1)\frac{\pi}{n})}{\sin(\frac{\pi}{n}) \sin(2\frac{\pi}{n}) \dots \sin(j\frac{\pi}{n})}.$$

This gives the x_j entry of $u_{>0}(t)$. To determine also the minors of $u_{>0}(t)$ and prove total positivity we will need a bit more notation.

Definition 8.3. Associate to a partition λ its Young diagram, and let the boxes in λ be labeled by their (x, y) coordinates in the usual way, as if they were entries of a matrix. We write $(i, j) \in \lambda$ if (i, j) are the coordinates of a box in λ , so $j \leq \lambda_i$ and $i \leq \lambda_j^t$. Let $\text{hl}(i, j)$ denote the *hook length* of the box labeled by (i, j) . That is, $\text{hl}(i, j) = \lambda_i + \lambda_j^t - i - j + 1$.

Theorem 8.4. *Recall the definitions of s_λ and q (from (5.1) and Remark 3.3).*

1. *For any partition λ , the value of s_λ on $u_{>0}(t)$ is given by the following hook length formula,*

$$s_\lambda(u_{>0}(t)) = t^{|\lambda|} \frac{\prod_{(i,j) \in \lambda} \sin((d-i+j)\frac{\pi}{n})}{\prod_{(i,j) \in \lambda} \sin(\text{hl}(i,j)\frac{\pi}{n})}.$$

The value of q is given by $q(u_{>0}(t)) = t^n$.

2. *The map $\mathbb{R} \xrightarrow{u_{>0}} \mathcal{V}_{d,n}(\mathbb{R})$ restricts to give homeomorphisms (for the standard Hausdorff topology) between*

$$\mathbb{R}_{\geq 0} \xrightarrow{\sim} \mathcal{V}_{d,n}(\mathbb{R}_{\geq 0}) \quad \text{and} \quad \mathbb{R}_{> 0} \xrightarrow{\sim} \mathcal{V}_{d,n}(\mathbb{R}_{> 0}).$$

So the $u_{>0}(t)$ for $t \in \mathbb{R}_{\geq 0}$ are precisely all the totally nonnegative elements in $\mathcal{V}_{d,n}$.

3. *An element $u \in \mathcal{V}_{d,n}$ lies in $\mathcal{V}_{d,n}(\mathbb{R}_{\geq 0})$ if and only if $s_\lambda(u) \in \mathbb{R}_{\geq 0}$, for all $\lambda \in \text{Sh}(d, c)$. It lies in $\mathcal{V}_{d,n}(\mathbb{R}_{> 0})$ precisely if $s_\lambda(u) \in \mathbb{R}_{> 0}$ for all $\lambda \in \text{Sh}(d, c)$.*

Proof. From the definitions it follows that $s_\lambda(u_{>0}(t)) = S_\lambda(t\zeta^{I_0}) = t^{|\lambda|} S_\lambda(\zeta^{I_0})$. Now the identity in (1) is equivalent to

$$S_\lambda(\zeta^{I_0}) = \frac{\prod_{(i,j) \in \lambda} \sin((d-i+j)\frac{\pi}{n})}{\prod_{(i,j) \in \lambda} \sin(\text{hl}(i,j)\frac{\pi}{n})}.$$

This equality is again the consequence of a classical formula. It is a variant of Littlewood's identity for $S_\lambda(1, q, q^2, \dots, q^k)$, see [12] Example I.3.1. The proof goes by writing the Schur polynomial (by Weyl's character formula) as quotient of two determinants. So $S_\lambda(x^{-\frac{d-1}{2}}, x^{-\frac{d-1}{2}+1}, \dots, x^{\frac{d-1}{2}}) = \frac{\Delta_\lambda^\lambda(x)}{\Delta_\emptyset^\emptyset(x)}$, where $\Delta_d^\lambda(x)$ is the generalized $d \times d$ Vandermonde determinant

$$\Delta_d^\lambda(x) = \det \left(x^{(-\frac{d-1}{2}+i-1)(\lambda_j+d-j)} \right) = \prod_{1 \leq i < j \leq d} \left(x^{\frac{\lambda_i - \lambda_j + i - j}{2}} - x^{\frac{\lambda_i - \lambda_j + j - i}{2}} \right).$$

Replacing x by ζ and $\left(x^{\frac{\lambda_i - \lambda_j + i - j}{2}} - x^{\frac{\lambda_i - \lambda_j + j - i}{2}} \right)$ by $2i \sin((\lambda_i - \lambda_j + i - j)\frac{\pi}{n})$ and cancelling the numerator against the denominator in S_λ the desired formula follows. The identity for q is immediate from Lemma 3.7.

(2) and (3). To prove that any $u \in U^+$ is totally nonnegative it suffices to check that all the minors with connected column sets are nonnegative, by a classical result. For $u = u_{>0}(t) = u_n(t\zeta^{I_0})$ these minors are precisely Schur polynomial values $S_\lambda(t\zeta^{-\frac{d-1}{2}}, t\zeta^{-\frac{d-1}{2}+1}, \dots, t\zeta^{\frac{d-1}{2}})$ where λ is a partition which fits inside a $j \times (n-j)$ box for some $j = 1, \dots, n-1$. In other words λ must be a partition with all hook lengths less than n . Therefore in the formula in (3) the denominator is always $\neq 0$ (if $t \neq 0$). From the numerator it follows that $s_\lambda(u_{>0}(t)) \neq 0$ precisely if $\lambda \in \text{Sh}(d, c)$.

Suppose now that $t > 0$ and $\lambda \in \text{Sh}(d, c)$. Then for any box (i, j) in λ both $d-i+j$ as well as the hook length are positive integers less than n . Therefore the values $\sin((d-i+j)\frac{\pi}{n})$ and $\sin(\text{hl}(i, j)\frac{\pi}{n})$ in the formula are all positive and thus these minors are > 0 . For $t = 0$ all S_λ except for $S_{(0, \dots, 0)} \equiv 1$ vanish. This implies (3) and one direction in (2).

It remains to prove that there are no other matrices in $\mathcal{V}_{d,n}(\mathbb{R}_{\geq 0})$. Suppose therefore $u = u_n(t\zeta^I)$ is totally nonnegative, where $t \in \mathbb{C}^*$ and $I \in \mathcal{I}_{d,n}$. We have observed that all of the s_λ for $\lambda \in \text{Sh}(d, c)$ must take nonnegative values on u .

Since $u_n(t\zeta^I) \in \mathcal{V}_{d,n}(\mathbb{R})$ we know that $q(u_n(t\zeta^I)) = t^n$ must be real. In fact, q can be expressed as

$$q = X_d Y_c$$

in $\Lambda_{d,n}$, which follows by expanding the determinant (3.1) for Y_n to

$$Y_n = \sum_{k=1}^{n-1} (-1)^{k+1} X_k Y_{n-k} = (-1)^{d+1} X_d Y_c.$$

Therefore $q(u(t\zeta^I)) = t^n$ must be positive real. So $t = \zeta^k r$ for some $k \in \{1, \dots, n\}$ and $r \in \mathbb{R}_{>0}$. We can replace t by r and ζ^I by $\zeta^J := \zeta^{I+(k, \dots, k)}$ to get $u_n(t\zeta^I) = u_n(r\zeta^J)$. If $\zeta^J = \zeta^{I_0}$ then $u_n(r\zeta^J) = u_{>0}(r)$, so we may assume that this is not the case. By the orthogonality formula (2) from Theorem 4.3 we then have,

$$\sum_{\lambda \in \text{Sh}(d,c)} S_\lambda(r\zeta^J) S_{\text{PD}(\lambda)}(r\zeta^{I_0}) = 0.$$

Since all the $S_{\text{PD}(\lambda)}(r\zeta^{I_0})$ are strictly positive, the $S_\lambda(r\zeta^J)$ cannot all be nonnegative. This finishes the proof. $\square$

9. FACTORIZATION OF ELEMENTS IN $\mathcal{V}_{d,n}(\mathbb{R}_{>0})$

The totally positive part $\mathcal{V}_{d,n}(\mathbb{R}_{>0})$ lies in particular in $U_{w_{\{\hat{d}\}}}^+(\mathbb{R}_{>0})$. Hence we can factorize its elements into products of elements $x_i(t) = \exp(te_i)$ (with $t \in \mathbb{R}_{>0}$) as described in (7.1).

Definition 9.1. For $k \leq d$ and $m \leq c$ let

$$B(k, m) := \{(i, j) \mid d - k + 1 \leq i \leq d, 1 \leq j \leq m\}.$$

Then $B(d, c)$ labels the set of boxes in the Young diagram of (c^d) and we may think of $B(k, m)$ as parameterizing a subset. We introduce a partial ordering $\preceq$ on $B(k, m)$ by

$$(i, j) \preceq (i', j') \iff i \geq i' \text{ and } j \geq j'.$$

There is a unique smallest element (d, m) and largest element $(d - k + 1, 1)$. When taking products over $B(k, m)$ we will consider the factors ordered compatibly with this partial order (in increasing order). So for example the product $\prod_{(i,j) \in B(d,c)} s_{d-i+j}$ of simple reflections in the Weyl group gives $w_{\{\hat{d}\}}$. The different allowed orderings of the factors correspond to different reduced expressions.

For each pair (k, m) we also have a well-defined monomial

$$e_{(k,m)} = \prod_{(i,j) \in B(k,m)} e_{d-i+j}$$

in the universal enveloping algebra $\mathcal{U}^+$ of $\text{Lie}(U^+)$.

Next we define

$$\begin{aligned} x : (\mathbb{C}^*)^{B(d,c)} &\rightarrow U^+ \cap B^- w_{\{\hat{d}\}} B^- \\ \mathbf{a} = (a_{(i,j)}) &\mapsto x(\mathbf{a}) := \prod_{(i,j) \in B(d,c)} x_{d-i+j}(a_{(i,j)}). \end{aligned}$$

So the restriction of x to $(\mathbb{R}_{>0})^{B(d,c)}$ is the parameterization map from (7.1).

Finally, let $\Delta_{[i,j]} \in \mathbb{C}[U^+]$ to be the minor with row set $1, \dots, 1 + j - i$ and column set $i, \dots, j$.

Lemma 9.2. *Let $u = x(\mathbf{a})$ for $\mathbf{a} = (a_{(i,j)}) \in (\mathbb{C}^*)^{B(d,c)}$.*

1. *The $m \times m$ -minor $\Delta_{[k+1,k+m]}$ of u is given by*

$$\Delta_{[k+1,k+m]}(u) = \prod_{(i,j) \in B(k,m)} a_{(i,j)}.$$

2. *The entries of $\mathbf{a}$ are computed in terms of the minors u by*

$$a_{(d-k+1,m)} = \frac{\Delta_{[k+1,k+m]}(u) \Delta_{[k,k+m-2]}(u)}{\Delta_{[k+1,k+m-1]}(u) \Delta_{[k,k+m-1]}(u)},$$

where $k \in \{1, \dots, d\}$ and $m \in \{1, \dots, c\}$.

The entries $a_{(i,j)}$ can also be worked out using Berenstein and Zelevinski's generalized Chamber Ansatz [1]. This method for factorizing elements of $U_w^+(\mathbb{R}_{>0})$ is more general in that it works for any w . But it gives the entries in the factorization of an element $x \in U_w^+(\mathbb{R}_{>0})$ in terms of minors not of x itself but of another matrix $z = \eta_w^{-1}(x)$, which in turn can be expressed using minors of x . In the case at hand it would also be possible to show that their map η_w is essentially the identity on these particular elements. This explains why the formula (2) looks like the one given by a single application of the Chamber Ansatz.

Proof. (1) Let $v_1, \dots, v_n$ be the standard basis of $\mathbb{C}^n$. Then the minor in question is just the matrix coefficient

$$(9.1) \quad \langle u \cdot v_{k+1} \wedge \dots \wedge v_n, v_1 \wedge \dots \wedge v_m \wedge v_{k+m+1} \wedge \dots \wedge v_n \rangle$$

in the fundamental representation $\bigwedge^{n-k} \mathbb{C}^n$ of $GL_n(\mathbb{C})$. Now consider the monomial $e(k, m) = \prod_{(i,j) \in B(k,m)} e_{d-i+j}$ in $\mathcal{U}^+$. It has the property that

$$e(k, m) \cdot v_{k+1} \wedge \dots \wedge v_n = v_1 \wedge \dots \wedge v_m \wedge v_{k+m+1} \wedge \dots \wedge v_n.$$

In fact, $e(k, m)$ is the unique (up to scalar) monomial in the e_i 's satisfying

$$\langle e(k, m) \cdot v_{k+1} \wedge \dots \wedge v_n, v_1 \wedge \dots \wedge v_m \wedge v_{k+m+1} \wedge \dots \wedge v_n \rangle \neq 0.$$

Then embed $u = \prod_{(i,j) \in B(d,c)} x_{d-i+j}^{a_{(i,j)}}$ into the (graded) completion of the enveloping algebra and expand and multiply the exponentials to get an infinite series of monomials in the e_i 's. The coefficient of $\prod_{(i,j) \in B(k,m)} e_{d-i+j}$ in this series is precisely $\prod_{(i,j) \in B(k,m)} a_{(i,j)}$. Since this is the only monomial that contributes to the matrix coefficient (9.1) the statement follows. Part (2) follows directly from (1) by cancellation. $\square$

Proposition 9.3. 1. *If $u = u_{>0}(t)$, then*

$$u = \prod_{(i,j) \in B(d,c)} x_{d-i+j} \left(t \frac{\sin((i+j-1)\frac{\pi}{n})}{\sin((d-i+j)\frac{\pi}{n})} \right),$$

where the order of multiplication is such that the indices of the x_{d-i+j} spell a reduced expression of $w^{\{\vec{d}\}}$.

2. *An element $u \in \mathcal{V}_{d,n}$ lies in $\mathcal{V}_{d,n}(\mathbb{R}_{>0})$ precisely if $s_{(m^k)}(u) > 0$ for all rectangular partitions (m^k) with $m \leq c$ and $k \leq d$.*

Proof. (1). The minor $\Delta_{[k+1, k+m]}$ of $u_n(t\zeta^I)$ is computed by $S_{(m^k)}(\zeta^I)$, by the dual Jacobi–Trudi formula. For $u_{>0}(t)$ this gives

$$\Delta_{[k+1, k+m]}(u_{>0}(t)) = S_{(m^k)}(t\zeta^{I_0}) = t^{km} \prod_{(i,j) \in (m^k)} \frac{\sin((d-i+j)\frac{\pi}{n})}{\sin((\text{hl}(i,j))\frac{\pi}{n})}.$$

Therefore $u_{>0}(t) = \prod_{(i,j) \in B(d,c)} x_{d-i+j}(a_{(i,j)})$ where $a_{(i,j)}$ is computed by

$$a_{(i,j)} = t \frac{\sin((i+j-1)\frac{\pi}{n})}{\sin((d-i+j)\frac{\pi}{n})}$$

after applying Lemma 9.2(2) and cancelling in the numerator and denominator.

(2) is clear since by (7.1) and 9.2 the totally positive part of $U^+ \cap B^{-w\{\hat{d}\}}B^-$ itself is determined by the inequalities $\Delta_{[m+1, \dots, m+k]} > 0$. In fact, this criterion is the earlier result of Berenstein and Zelevinski [1], which we have just reproved in our special case. Now because $\Delta_{[m+1, \dots, m+k]}|_{\mathcal{V}_{d,n}} = s_{(m^k)}$, we are done. $\square$

Note that the reduced set of cd inequalities given in Proposition 9.3(2) to describe $\mathcal{V}_{d,n}(\mathbb{R}_{>0})$ inside variety $\mathcal{V}_{d,n}(\mathbb{R})$ is still surprisingly large, for a 1-dimensional variety. It is the same as the set needed to describe the positive part $U_{w\{\hat{d}\}}(\mathbb{R}_{>0})$ in the whole cd -dimensional variety $U_{w\{\hat{d}\}}$.

10. QUANTUM COHOMOLOGY OF GRASSMANNIANS

This section is a brief collecting together of some basic facts about the (small) quantum cohomology ring of a Grassmannian. Quantum cohomology originally comes from the work of Vafa and Witten and ideas from string theory. Since then it has had a big impact on mathematics, with much work being done to make the theory rigorous and expand on it. Our main reference is Bertram’s self-contained mathematical treatment for the Grassmannian [2]. See also [5] and [13] and references therein for a more general introduction to the mathematical theory of quantum cohomology (in the algebraic geometric or symplectic setting respectively).

Let $Gr_d(n)$ denote the variety of all d -dimensional subspaces of $\mathbb{C}^n$. As before we fix $c = n - d$. So c is the codimension while d is the dimension.

10.1. Schubert basis. The quantum cohomology of $Gr_d(n)$ is in the first instance a module over the polynomial ring in one variable $\mathbb{Z}[q]$ defined by

$$(10.1) \quad qH^*(Gr_d(n)) = \mathbb{Z}[q] \otimes_{\mathbb{Z}} H^*(Gr_d(n), \mathbb{Z}).$$

Therefore it has a $\mathbb{Z}[q]$ -basis given by the classical Schubert basis of $H^*(Gr_d(n), \mathbb{Z})$. We briefly recall its definition here.

Consider a flag $V_{\bullet}$ of subspaces $V_{\bullet} = (V_1 \subset V_2 \subset \dots \subset V_n = \mathbb{C}^n)$ and a partition $\lambda \in \text{Sh}(d, c)$. Then the Schubert variety $X_{\lambda}(V_{\bullet}) \subset Gr_d(n)$ is defined concretely as

$$X_{\lambda}(V_{\bullet}) = \{W \in Gr_d(n) \mid \dim(W \cap V_{c+i-\lambda_i}) \geq i\}.$$

$X_{\lambda}(V_{\bullet})$ is a closed subvariety of $Gr_d(n)$ of (complex) codimension $|\lambda|$. Since any flag can be transformed into any other by the action of $GL_n(\mathbb{C})$, the homology class of $X_{\lambda}(V_{\bullet})$ does not depend on the choice of flag $V_{\bullet}$. By the classical result of Ehresmann, the Schubert varieties give rise to a basis of the homology of $Gr_d(n)$. Let us denote the Poincaré dual basis of cohomology by $\{\sigma_{\lambda} \mid \lambda \in \text{Sh}(d, c)\}$. So we have $\sigma_{\lambda} \in H^{2|\lambda|}(Gr_d(n))$.

10.2. Poincaré duality. Consider the Poincaré duality pairing

$$(\ , \) : H^*(Gr_d(n)) \times H^*(Gr_d(n)) \rightarrow \mathbb{Z}$$

defined by $(\sigma, \tau) = (\sigma \cup \tau)[Gr_d(n)]$, where $[Gr_d(n)] \in H_{2cd}(Gr_d(n))$ is the fundamental class. We recall that in terms of the Schubert basis elements this pairing takes the form

$$(10.2) \quad (\sigma_\lambda, \sigma_\mu) = \delta_{\lambda, PD(\mu)},$$

where as before $PD(\mu) = (c - \mu_d, c - \mu_{d-1}, \dots, c - \mu_1)$.

10.3. Ring structure. As a ring, $qH^*(Gr_d(n))$ does not agree with $H^*(Gr_d(n)) \otimes_{\mathbb{Z}} \mathbb{Z}[q]$. The new ring structure is a q -deformation of the old with structure constants for the Schubert basis given by 3-point Gromov–Witten invariants (see below). Let us view our Schubert basis elements $\{\sigma_\lambda \mid \lambda \in \text{Sh}(d, c)\}$ as lying inside $qH^*(Gr_d(n))$ by identifying σ_λ with $\sigma_\lambda \otimes 1$.

Let $\lambda, \mu, \nu \in \text{Sh}(d, c)$ such that $|\lambda| + |\mu| + |\nu| = dc + kn$ (the dimension of the moduli space of degree k holomorphic maps $\mathbb{CP}^1 \rightarrow Gr_d(n)$), and fix three flags $V_\bullet, V'_\bullet$ and $V''_\bullet$ in general position. Then $\langle \sigma_\lambda, \sigma_\mu, \sigma_\nu \rangle_k$ is defined as the number of degree k holomorphic curves $\mathbb{CP}^1 \rightarrow Gr_d(n)$ taking $0, 1$, and $\infty \in \mathbb{CP}^1$ to points in $X_\lambda(V_\bullet)$, $X_\mu(V'_\bullet)$ and $X_\nu(V''_\bullet)$, respectively. If $|\lambda| + |\mu| + |\nu| \neq dc + kn$ then $\langle \sigma_\lambda, \sigma_\mu, \sigma_\nu \rangle_k := 0$.

Multiplication in $qH^*(Gr_d(n))$ is defined by

$$\sigma_\lambda \cdot \sigma_\mu = \sum_{\nu, k} \langle \sigma_\lambda, \sigma_\mu, \sigma_{PD(\nu)} \rangle_k q^k \sigma_\nu.$$

It is a remarkable fact proved now in more general contexts by Ruan–Tian [18] and also Kontsevich–Manin [7] that the quantum cohomology product is associative. For the Grassmannian the associativity of the above defined multiplication is also derived directly in [2]. With this product $qH^*(Gr_d(n))$ becomes a graded ring with grading defined by $\deg(\sigma_\lambda) = |\lambda|$ and $\deg(q) = n$ (or everything doubled to agree with the grading of the ordinary cohomology). Note that the structure constants of this ring are nonnegative integers. Also when q is set to zero only the $k = 0$ terms in the product formula remain, and we recover the definition of the usual cup-product.

10.4. Presentation and structure constants. Recall the definition of the ring $\Lambda_{d,n}^{\mathbb{Z}}$ from Definition 3.2 (and Corollary 5.2).

Theorem 10.1 ([19], [20]). *The ring $qH^*(Gr_d(n))$ is generated by the d Schubert classes $\sigma_{(1)}, \sigma_{(1,1)}, \dots, \sigma_{(1^d)}$. The assignment*

$$(10.3) \quad \Lambda_{d,n}^{\mathbb{Z}} \rightarrow qH^*(Gr_d(n)) : \quad X_k \mapsto \sigma_{(1^k)}, \quad (-1)^{d+1} Y_n \mapsto q$$

defines an isomorphism of rings.

It is also known that, as in the classical cohomology, the determinants $Y_1, \dots, Y_c$ map to the Schubert classes $\sigma_{(1,0,\dots,0)}, \dots, \sigma_{(c,0,\dots,0)}$. In fact, Bertram proved that all the classical Giambelli formulas hold also in the quantum cohomology ring.

Theorem 10.2 (quantum Giambelli formula, [2]). *For $\lambda \in \text{Sh}(d, c)$, the element $s_\lambda \in \Lambda_{d,n}^{\mathbb{Z}}$ defined in (5.1) is mapped to the (quantum) Schubert class σ_λ under the isomorphism (10.3).*

With this theorem we recover the Bertram–Vafa–Intriligator formula for the Gromov–Witten invariants from our algebraic result Corollary 6.2.

Theorem 10.3 ([2], Section 5). *The Gromov–Witten invariant $\langle \sigma_\lambda, \sigma_\mu, \sigma_\nu \rangle_k$ where $|\lambda| + |\mu| + |\nu| = nk + cd$ is computed by*

$$\langle \sigma_\lambda, \sigma_\mu, \sigma_\nu \rangle_k = \frac{1}{n^d} \sum_{J \in \mathcal{I}_{d,n}} S_\lambda(\zeta^J) S_\mu(\zeta^J) S_\nu(\zeta^J) \frac{|\text{Vand}(\zeta^J)|^2}{S_{(cd)}(\zeta^J)}$$

□

See also [19] for a more general version the Vafa–Intriligator formula.

10.5. Poincaré Duality for $qH^*(Gr_d(n))$. We end this section with the geometric interpretation of the ‘row–orthogonality’ formula (4.7). One can define an analogue of the Poincaré duality pairing on $qH^*(Gr_d(n))$. This is a $\mathbb{Z}[q]$ –bilinear map

$$(\ , \)_q : qH^*(Gr_d(n)) \times qH^*(Gr_d(n)) \rightarrow \mathbb{Z}[q]$$

which takes (σ, σ') to the coefficient of $\sigma_{(cd)}$ in $\sigma \cdot \sigma'$. The pairing $(\ , \)_q$ specializes to usual Poincaré duality when q goes to 0. For two Schubert basis elements σ_λ and σ_μ this pairing is simply computed by

$$(10.4) \quad (\sigma_\lambda, \sigma_\mu)_q = \langle \sigma_\lambda, \sigma_\mu, \sigma_{PD(cd)} \rangle_k q^k$$

if n divides $|\lambda| + |\mu| - cd$ with quotient k . And it is zero if $|\lambda| + |\mu| - cd$ is not divisible by n . Now by Theorem 10.3, equation (10.4) becomes

$$(\sigma_\lambda, \sigma_\mu)_q = \frac{1}{n^d} \sum_{J \in \mathcal{I}_{d,n}} S_\lambda(\zeta^J) S_\mu(\zeta^J) \frac{|\text{Vand}(\zeta^J)|^2}{S_{(cd)}(\zeta^J)}.$$

Thus the equation (4.7) corresponds to the following geometric statement.

Proposition 10.4 ([20] Section 3.1, [2] Lemma 2.5.). *Let $\lambda, \mu \in \text{Sh}(d, c)$. Then the pairing $(\ , \)_q$ defined above takes the same form*

$$(\sigma_\lambda, \sigma_\mu)_q = \delta_{\lambda, PD(\mu)}$$

as the classical Poincaré duality pairing.

□

11. AN INEQUALITY FOR SCHUR POLYNOMIAL VALUES

In this section we give a little application of the positivity of structure constants in $\Lambda_{d,n}$. The positivity of structure constants is one property of $\Lambda_{d,n}$ coming from its isomorphism with $qH^*(Gr_d(n), \mathbb{C})$ for which we know no elementary explanation. The Vafa–Intriligator formula does not obviously give something positive. And there is a generalization of the Littlewood–Richardson rule to quantum cohomology in [3], but it is an alternating formula.

Proposition 11.1. *Let $I_0 = (-\frac{d-1}{2}, \dots, \frac{d-1}{2})$ and $I \in \mathcal{I}_{d,n}$ any other element, and let $\zeta = \exp(\frac{2\pi i}{n})$ as before. Then for any partition $\lambda \in \text{Sh}(d, c)$,*

$$(11.1) \quad |S_\lambda(\zeta^I)| \leq S_\lambda(\zeta^{I_0}).$$

Remark 11.2. An equivalent formulation of the above result is to say that

$$|S_\lambda(\zeta^{i_1}, \dots, \zeta^{i_d})| \leq S_\lambda(\zeta^{-\frac{d-1}{2}}, \zeta^{-\frac{d-1}{2}+1}, \dots, \zeta^{\frac{d-1}{2}})$$

for any d –tuple of distinct n –th roots of unity $\zeta^{i_1}, \dots, \zeta^{i_d}$ and partition $\lambda \in \text{Sh}(d, c)$.

Proof. Consider $qH^*(Gr_d(n), \mathbb{C})_{q=1} := qH^*(Gr_d(n), \mathbb{C}) \otimes_{\mathbb{C}[q]} \mathbb{C}[q]/(q-1)$, the specialization of the quantum cohomology ring at $q = 1$. We may identify this ring with the ring of functions on the fiber of $\mathcal{V}_{d,n}$ over $q = 1$, that is with functions on the finite set $\{u_n(\zeta^I) \mid I \in \mathcal{I}_{d,n}\}$. We have that $qH^*(Gr_d(n), \mathbb{C})_{q=1}$ is a finite dimensional vector space with basis given by (restriction of) the Schubert basis σ_λ .

Another basis of this space is the set consisting of all $\sigma_I := \sum_{\nu \in \text{Sh}(d,c)} \overline{S_\nu(\zeta^I)} \sigma_\nu$ for $I \in \mathcal{I}_{d,n}$. By Proposition 4.3 (3) we have

$$\sigma_I(u_n(\zeta^J)) = \delta_{I,J} \frac{n^d}{|\text{Vand}(\zeta^I)|^2},$$

so these are up to scalar the characteristic functions of the points in the set $\{u_n(\zeta^I) \mid I \in \mathcal{I}_{d,n}\}$.

Now consider the multiplication operator $[\sigma_\lambda] : \sigma \mapsto \sigma_\lambda \sigma$ on $qH^*(Gr_d(n), \mathbb{C})_{q=1}$. Then clearly

$$\sigma_\lambda \cdot \sigma_I = S_\lambda(\zeta^I) \sigma_I.$$

So σ_I is an eigenvector of the multiplication operator $[\sigma_\lambda]$ with eigenvalue $S_\lambda(\zeta^I)$. Moreover the set $\{\sigma_I \mid I \in \mathcal{I}_{d,n}\}$ is a complete eigenbasis, in fact a simultaneous eigenbasis for all the multiplication operators $[\sigma]$, where $\sigma \in qH^*(Gr_d(n), \mathbb{C})_{q=1}$.

Consider the set Y of all $\sigma \in qH^*(Gr_d(n), \mathbb{R})_{(q=1)}$ such that all the eigenvalues of $[\sigma]$ are distinct. Then Y is obtained from the real vector space $qH^*(Gr_d(n), \mathbb{R})_{(q=1)}$ by removing certain lower dimensional linear subspaces. Therefore in particular Y is open dense in $qH^*(Gr_d(n), \mathbb{R})_{(q=1)}$.

Now recall that the multiplication operators $[\sigma_\lambda]$ on $qH^*(Gr_d(n), \mathbb{C})_{q=1}$ are given in terms of the Schubert basis by an $\binom{n}{d} \times \binom{n}{d}$ -matrix A_λ with nonnegative integer entries (Gromov–Witten invariants). We can approximate the matrix A_λ to arbitrary precision $\epsilon > 0$ by some other nonnegative matrix

$$A_\lambda^\epsilon := A_\lambda + \sum_{\mu \in \text{Sh}(d,c)} \epsilon_\mu A_\mu, \quad \text{where } 0 < \epsilon_\mu < \epsilon,$$

which has only simple eigenvalues, by the previous paragraph. We now apply the following version of the Perron–Frobenius eigenvalue theorem to A_λ^ϵ .

Perron–Frobenius Theorem. ([14] I. Theorem 4.2) Suppose A is an $m \times m$ matrix with all entries in $\mathbb{R}_{\geq 0}$. Then A has an eigenvalue r in $\mathbb{R}_{\geq 0}$ with a nonnegative eigenvector and such that all other eigenvalues of A have absolute value $\leq r$.

Let $A = A_\lambda^\epsilon$ chosen as above. Since all of its eigenvalues are simple, the eigenvectors σ_I are unique up to scalar. Therefore using Theorem 8.4 it follows that (up to positive scalar) σ_{I_0} is the only nonnegative eigenvector. So by the Perron–Frobenius theorem its eigenvalue, $S_\lambda(\zeta^{I_0}) + \sum_\mu \epsilon_\mu S_\mu(\zeta^{I_0})$ is the maximal eigenvalue. Now choosing ϵ sufficiently small we obtain that $S_\lambda(\zeta^{I_0})$ must be a maximal eigenvalue of A_λ . This proves the inequality (11.1). $\square$

REFERENCES

1. A. Berenstein and A. Zelevinsky, *Total positivity in Schubert varieties*, Comment. Math. Helv. **72** (1997), 128–166.
2. A. Bertram, *Quantum Schubert calculus*, Advances in Mathematics **128** (1997), 289–305.
3. A. Bertram, I. Ciocan-Fontanine, and W. Fulton, *Quantum multiplication of Schur polynomials*, J. Algebra **219** (1999), no. 2, 728–746.
4. W. Fulton, , L.M.S. Student Texts, no. 35, Cambridge Univ. Press, 1997.

5. W. Fulton and R. Pandharipande, *Notes on stable maps and quantum cohomology*, Algebraic geometry—Santa Cruz 1995 (Providence, RI), Amer. Math. Soc., Providence, RI, 1997, pp. 45–96.
6. D. Gepner, *Fusion rings and geometry*, Comm. Math. Phys. **141** (1991), 381–411.
7. M. Kontsevich and Yu. Manin, *Gromov-Witten classes, quantum cohomology, and enumerative geometry*, Comm. Math. Phys. **164** (1994), 525–562.
8. B. Kostant, *Flag manifold quantum cohomology, the Toda lattice, and the representation with highest weight ρ* , Selecta Math. (N.S.) **2** (1996), 43–91.
9. G. Lusztig, , Progress in Mathematics, Birkhaeuser, Boston, 1993.
10. ———, *Total positivity in reductive groups*, Lie theory and geometry: in honor of Bertram Kostant (Boston) (G. I. Lehrer, ed.), Progress in Mathematics, vol. 123, Birkhaeuser, Boston, 1994, pp. 531–568.
11. ———, *Total positivity in partial flag manifolds*, Representation Theory **2** (1998), 70–78.
12. I. G. Macdonald, *Symmetric functions and Hall polynomials*, 2nd ed., Oxford Univ. Press, 1995.
13. Dusa McDuff and Dietmar Salamon, , University Lecture Series, American Mathematical Society, Providence, RI, 1994.
14. Henryk Minc, *Nonnegative matrices*, John Wiley & Sons Inc., New York, 1988.
15. D. Peterson, *Quantum cohomology of G/P* , Lecture Course, spring term, M.I.T., 1997.
16. ———, *Quantum cohomology of G/P* , Seminaire de Mathematiques Superieures: Representation Theories and Algebraic Geometry, Universite de Montreal, Canada, July 28-Aug. 8 1997 (unpublished lecture notes).
17. K. Rietsch, *Quantum cohomology of partial flag varieties and totally positive Toeplitz matrices*, preprint.
18. Y. Ruan and G. Tian, *A mathematical theory of quantum cohomology*, J. Diff. Geom. **42** (1995), 259–367.
19. B. Siebert and G. Tian, *On quantum cohomology rings of Fano manifolds and a formula of Vafa and Intriligator*, Asian J. Math. **1** (1997), no. 4, 679–695.
20. E. Witten, *The Verlinde algebra and cohomology of the Grassmannian*, Geometry, topology & Physics, Conf. Proc. Lecture Notes **VI** (1995), 357–422.

DPMMS, CAMBRIDGE UNIVERSITY AND NEWNHAM COLLEGE, CAMBRIDGE
E-mail address: `rietsch@dpmms.cam.ac.uk`